

# Sustainability of Home Telepathy Programs: A Systematic Review



Kavita Radhakrishnan<sup>1</sup>, Bo Xie<sup>1</sup>, Amy Ellis<sup>1</sup>

As part of the Medicare Care Transitions Act of 2009, the federal government mandated reductions in re-hospitalizations through better care coordination and follow up services. Remote monitoring technologies such as telehealth has emerged as a potential solution to reduce re-hospitalization and healthcare utilization costs and manage chronic diseases in the home health community. However, sustainability of home telehealth programs remains a major challenge with unclear understanding of factors contributing to discontinued or sustained telehealth use. Earlier systematic reviews have focused on the effectiveness of home telehealth programs for physiological or behavioral outcomes,<sup>1-4</sup> but they have not addressed such programs' sustainability.

**Methods:** To address this knowledge gap, we present here a systematic review of articles published from 1996 to 2013 within the databases of CINAHL, Pubmed/Medline, PsychInfo, Web of Science, Cochrane Reviews to identify barriers to and facilitators for sustained telehealth use by home health patients and clinicians for chronic physiological disease management. For this review, we used the search terms of telehealth, telemonitoring, telecare, telemedicine, and telehomecare and adapted Craddock's definition of sustainability for telehealth services as the use of home telehealth services that holds the promise of being absorbed into routine health-care delivery including an increasing demand for those services, as well as acceptance of such services among healthcare providers along with a commitment to invest in them. Articles were included if they reported on longitudinal investigations of telehealth usage by home health agencies and addressed the management of chronic cardiovascular disease, diabetes, and obstructive pulmonary disease in older adults age 65 years or above. Data extraction using PRISMA guidelines and quality appraisal using Mixed Methods Appraisal Tool (MMAT) was conducted on relevant empirical studies. Thematic analysis across the studies and narrative summaries were used to synthesize the findings from the included studies. From the final articles, the following data were extracted: (1) study design; (2) study quality; (3) characteristics of the participants, including demographics, diagnoses, and role in the telehealth program; (4) data collection methods; (5) description of the telehealth program model, and (6) determinants of the sustainability of home telehealth programs.

**Results:** The initial 3920 citations were reduced to 943 after applying the initial search criteria and eliminating duplicates. After title and abstract search, we abstracted 142 full articles of which 18 articles,<sup>5-22</sup> of moderate quality met the inclusion criteria. Full-texts were retrieved by a graduate research assistant and reviewed by the first two authors. Majority of the studies were conducted in UK (9) & US (7), with 1 in Canada and the Netherlands each. The articles are recent; 12 of the 18 studies were published after 2010. Twelve of the studies had qualitative designs; these included case study, phenomenological, and ethnographic approaches, as well as 3 process evaluations of randomized controlled trials. The other studies included five quantitative studies which included 1 descriptive usability study, 1 survey and 3 secondary analyses of retrospective data; and 1 mixed methods study. Sample sizes ranged from 12 to 82 for the qualitative studies and from 132 to 403 for the survey and secondary analysis studies. Participants included only patients (10 studies), only clinicians (4), or mixed samples of both patients and clinicians (4). Patient diagnoses targeted by the telehealth programs included only heart failure (5 studies), only COPD (5), only diabetes (2), or any of those three chronic diseases (7). Major themes that on sustainability of home telehealth programs included: user perceptions on effectiveness of home telehealth programs for achieving intended outcomes, tailoring of home telehealth programs to patient characteristics and needs, communication and collaboration among telehealth users, home health organizational processes and culture, and technology usability and innovation.

**Discussion:** In summary, to realize the potential of telehealth services for chronic disease management, future program redesign must (1) recognize formal reorganization of work between the staffs of home health service settings to include partnership and accountability negotiation, system interoperability, and shared visions for patient care management; (2) identify criteria for patient characteristics to enable telehealth service delivery tailored to individual patients' capabilities and context; (3) include clear guidelines and protocols for patient teaching, mechanisms for feedback and response, and negotiation of patient responsibilities, empowering patients to become self-reliant in their care management; (4) include stakeholder input during program implementation for improved incorporation within workflow and life routines; (5) improve technical quality of communication; and (6) enhance device usability tailored to elder use.

**Conclusion:** The findings of this systematic review have important implications for sustained usage of telehealth programs by home health service settings and can help realize the potential of telehealth for chronic disease management.

<sup>1</sup>University of Texas

## References

1. Klersy, C, De Silvestri, A, Gabutti, G, Regoli, F, and Auricchio, A. A meta-analysis of remote monitoring of heart failure patients. *Journal of the American College of Cardiology* 2009;54:1683–1694.
2. Paré, G, Moqadem, K, Pineau, G, and St-Hilaire, C. Clinical effects of home telemonitoring in the context of diabetes, asthma, heart failure and hypertension: a systematic review. *Journal of Medical Internet Research* 2010;12:e21.
3. Radhakrishnan, K and Jacelon, C. Impact of telehealth on patient self-management of heart failure: a review of literature. *The Journal of Cardiovascular Nursing* 2012;27:33–43.
4. Udsen, FW, Hejlesen, O, and Ehlers, LH. A systematic review of the cost and cost-effectiveness of telehealth for patients suffering from chronic obstructive pulmonary disease. *Journal of Telemedicine and Telecare* 2014;20:212–220.
5. Fairbrother, P, Pinnock, H, Hanley, J, McCloughan, L, Sheikh, A, Pagliari, C, McKinstry, B, and TELESCOT programme team. Continuity, but at what cost? The impact of telemonitoring COPD on continuities of care: a qualitative study. *Primary Care Respiratory Journal: Journal of the General Practice Airways Group* 2012;21:322–328.
6. Fairbrother, P, Ure, J, Hanley, J, McCloughan, L, Denvir, M, Sheikh, A, McKinstry, B, and Telescot programme team. Telemonitoring for chronic heart failure: the views of patients and healthcare professionals - a qualitative study. *Journal of Clinical Nursing* 2014;23:132–144.
7. Gale, N and Sultan, H. Telehealth as 'peace of mind': embodiment, emotions and the home as the primary health space for people with chronic obstructive pulmonary disorder. *Health & Place* 2013;21:140–147.
8. Guzman-Clark, JRS, Servellen, G van, Chang, B, Menten, J, and Hahn, TJ. Predictors and outcomes of early adherence to the use of a home telehealth device by older veterans with heart failure. *Telemedicine Journal and E-Health: The Official Journal of the American Telemedicine Association* 2013;19:217–223.
9. Hardisty, AR, Peirce, SC, Preece, A, et al. Bridging two translation gaps: a new informatics research agenda for telemonitoring of chronic disease. *International Journal of Medical Informatics* 2011;80:734–744.
10. Hibbert, D, Mair, FS, May, CR, Boland, A, O'Connor, J, Capewell, S, and Angus, RM. Health professionals' responses to the introduction of a home telehealth service. *Journal of Telemedicine and Telecare* 2004;10:226–230.
11. Horton, K. The use of telecare for people with chronic obstructive pulmonary disease: implications for management. *Journal of Nursing Management* 2008;16:173–180.
12. Juretic, M, Hill, R, Hicken, B, Luptak, M, Rupper, R, and Bair, B. Predictors of attrition in older users of a home-based monitoring and health information delivery system. *Telemedicine Journal and E-Health: The Official Journal of the American Telemedicine Association* 2012;18:709–712.
13. Kaufman, DR, Pevzner, J, Hilliman, C, Weinstock, RS, Teresi, J, Shea, S, and Starren, J. Redesigning a Telehealth Diabetes Management Program for a Digital Divide Seniors Population. *Home Health Care Management & Practice* 2006;18:223–234.
14. LaFramboise, LM, Woster, J, Yager, A, and Yates, BC. A technological life buoy: patient perceptions of the Health Buddy. *The Journal of Cardiovascular Nursing* 2009;24:216–224.
15. Lamothe, L, Fortin, JP, Labbé, F, Gagnon, MP, and Messikh, D. Impacts of telehomecare on patients, providers, and organizations. *Telemedicine Journal and E-Health: The Official Journal of the American Telemedicine Association* 2006;12:363–369.
16. Mair, FS, Hiscock, J, and Beaton, SC. Understanding factors that inhibit or promote the utilization of telecare in chronic lung disease. *Chronic Illness* 2008;4:110–117.
17. Peeters, JM, Veer, AJE de, Hoek, L van der, and Francke, AL. Factors influencing the adoption of home telecare by elderly or chronically ill people: a national survey. *Journal of Clinical Nursing* 2012;21:3183–3193.
18. Radhakrishnan, K, Jacelon, CS, Bigelow, C, Roche, JP, Marquard, JL, and Bowles, KH. Association of comorbidities with home care service utilization of patients with heart failure while receiving telehealth. *The Journal of Cardiovascular Nursing* 2013;28:216–227.
19. Radhakrishnan, K, Jacelon, C, and Roche, J. Perceptions on the Use of Telehealth by Homecare Nurses and Patients With Heart Failure A Mixed Method Study. *Home Health Care Management & Practice* 2012;24:175–181.
20. Rogers, A, Kirk, S, Gately, C, May, CR, and Finch, T. Established users and the making of telecare work in long term condition management: implications for health policy. *Social Science & Medicine (1982)* 2011;72:1077–1084.
21. Sandberg, J, Trief, PM, Izquierdo, R, et al. A qualitative study of the experiences and satisfaction of direct telemedicine providers in diabetes case management. *Telemedicine Journal and E-Health: The Official Journal of the American Telemedicine Association* 2009;15:742–750.